

Formulation and *in vitro* Evaluation of Topical Liposomal Gel of Acitrecin and Methotrexate

Dr.Jithendra Patel,Mohammed salman

Avanathi Institute of Pharmaceutical Sciences,Hyderabad,Telangana,India.

ABSTRACT

Acitrecin and Methotrexate used in treatment of skin inflammatory diseases. liposomes can increase the deposition of drugs within the skin at the site of action and reduce side effects of drug. The aim of the present study was to formulate and evaluate liposomal vesicles loaded with Acitrecin and Methotrexate. liposomes containing Acitrecin and Methotrexate were prepared using thin film method. The quantities of lecithin and cholesterol were changed to enhance the encapsulation of the drug. Carbomer 940 was used as gel base and four different gel formulations including hydroalcoholic gel, MLV liposomal gel, SUV liposomal gel and blank MLV gel containing free drug were prepared. The release profile of Acitrecin and Methotrexate was determined using dialysis membrane method. Liposomes were also characterized by optical microscope and their particle size was also determined. Formulation containing lecithin: drug: cholesterol (100: 10: 10) having about 90.05 percent encapsulation was selected as the best formulation and the results of release showed SUV liposomal gel has the most regular and the least interaction between the drug and polymer. Results of particle size determination showed 50% of MLV and SUV liposomes had diameter below 33.80 μm and 22.09 μm , respectively. The results of characterization of the vesicles indicated the potential application of Acitrecin and Methotrexate loaded liposome as carrier system.

Key words: *Acitrecin and Methotrexate, liposome, thin film, gel*

INTRODUCTION

Transdermal drug delivery has many advantages over the other traditional routes of administration. But, the barrier nature of the skin is a significant obstacle for most drugs to be delivered into and through it. Many researchers have been investigated the liposomes as drug delivery systems that can improve transdermal drug delivery. Liposomes are spherical vesicles with concentric phospholipid bilayers [1]. The results of some studies have been indicated that liposomes can increase the deposition of drugs within the skin at the site of action and reduce its absorption into blood [2, 3]. Lipid bilayered membrane encloses an aqueous core that hydrophilic drugs get entrapped in the central aqueous core of the vesicles while lipophilic drugs get entrapped within the bilayered membrane [4]. Drugs encapsulated in liposomes can be transported without rapid degradation and minimum side effects to the recipients [5]. Liposomes have many advantages such as, biocompatibility and biodegradability, prolonging release of active pharmaceutical agents, ability to entrap lipophilic and hydrophilic drugs, protecting encapsulated agents from metabolic processes, nontoxicity, penetration enhancing and slow drug release [6-10].

Acitrecin and Methotrexate is a glucocorticoid that it is used in treatment of allergic reactions, asthma, rhinitis and other inflammatory diseases. Mechanism of action of the drug is control of the synthesis of prostaglandins and leukotrienes [11].

The aim of the present study was to prepare and evaluate topical liposome gel formulation of Acitrecin and Methotrexate and to study their *in vitro* release behavior.

MATERIALS AND METHODS

Egg lecithin and vitamin E (alpha tocopherol) were purchased from Merck, Germany. Cholesterol and Acitrecin and Methotrexate were obtained from BDH, England and Sicro, Italy respectively. Carbomer 940 was purchased from Antwerpen, Belgium. All of the solvents were of analytical grade.

Experimental Preparation of liposome

Liposomes were prepared by thin film method. Briefly egg lecithin, cholesterol, vitamin E (as antioxidant) and 250 mg Acitrecin and 500 mg Methotrexate were dissolved in 5ml chloroform. The quantities of lecithin and cholesterol were changed to enhance loading drug in liposomes (table 1). Then the mixture was evaporated in a rotary evaporator (Heidolph, WB2000, Germany) at 150 rpm for 30 min. The thin film formed in the round-bottomed flask was hydrated with phosphate buffer saline for 30 min. Glass ball was used for dispersion of liposome [1, 10].

Table 1: Effect of lipid composition on encapsulation efficiency of Acitrecin and Methotrexate liposomes

Formulation NO.	Compositions ratio (mg) (lecithin:drug:cholesterol)	Entrapment efficiency (%)	Formulation NO.	Compositions ratio (mg) (lecithin:drug:cholesterol)	Entrapment efficiency (%)
1	100:10:5	75.80	7	125:10:10	58.83
2	100:10:10	90.05	8	125:10:20	47.85
3	100:10:20	78.71	9	150:10:10	47.97
4	100:10:40	53.19	10	150:10:20	44.90
5	100:10:60	69.49	11	200:10:10	48.53
6	100:10:80	67.19	12	200:10:20	45.35

Preparation of SUV liposome

SUV liposomes were obtained by sonication of MLV liposome in an ultrasonic bath (F, Germany) 30 min at room temperature.

Determination of Acitrecin and Methotrexate entrapment in liposomes

The liposomal suspension was centrifuged at 20000 rpm for 30 min. The supernatant was removed and the liposomes were disrupted with ethanol 70% and the quantity of drug was measured using a spectrophotometer (Biotek Kontron, 922, England) at 239 nm.

Particle size determination

The average diameter of liposomes was determined using a particle sizer (Malvern, Xlongbed ver.2.15, England) at 25 °C.

Microscopy

liposomes were examined by optical microscope (Olympus, R4, Japan) to determine the shape and lamellarity of vesicles (magnification x 200).

Preparation of liposomal gel

Carbomer 940 (1%) was added slowly to a PBS buffer solution (pH 7.4), under constant stirring by a paddle stirrer. For gel preservation, methyl paraben (0.2%) was added. Then triethanolamine was added for achieving neutral pH and clearing of the gels. After addition of the full amount of solid material, the gels were

allowed to swell under moderate stirring. Four different formulation gel including hydro-alcoholic gel, MLV liposomal gel, SUV liposomal gel and blank MLV gel plus the drug were prepared (table 2).

Table 2: Different gel formulations of Acitrecin and Methotrexate.

F ₁	hydroalcoholic gel
F ₂	MLV liposomal gel (containing lecithin 100 mg: drug 10 mg: cholesterol: 10 mg)
F ₃	SUV liposomal gel (containing lecithin 100 mg: drug 10 mg: cholesterol: 10 mg)
F ₄	blank MLV liposome gel plus the drug

Stability evaluation of liposomal gels

The liposomal formulations were stored in refrigerator for 3 months to evaluate color and appearance. 1 gr of each sample was dispersed in 100 ml distilled water and pH was determined using a pH meter (Coring, 7, England).

In vitro drug release studies

In vitro Acitrecin and Methotrexate release from the liposomes was measured using dialysis membrane method. 1 g of each formulation was put in a dialysis bag (Medicell International England). The receptor phase was contained ethanol 20% and phosphate buffer saline (pH 7.4) was stirred continually using a magnet stirrer at 37°C. An aliquot of 2.5 ml of samples were withdrawn from each batch at definite time intervals (15, 30, 45, 60, 90, 120, 150, 180, 210, 240, 270 and 300 min) and replaced with the same amount of ethanol and phosphate buffer saline to maintain sink condition. Then, the concentration of released drug was monitored using a UV spectrophotometer at 239 nm.

Statistical analysis of data

The reported release data were analyzed by Jim fit program [12] and compared and plotted using Excel Microsoft. $P > 0.05$ was considered as significant difference.

RESULTS AND DISCUSSION

As mentioned before, for determination of entrapment efficiency, the quantities of lecithin and cholesterol were changed to enhance loading drug in liposomes. At first, the quantity of cholesterol was changed (formulation No.1-6). Then the quantities of cholesterol (10 and 20 mg) that had high encapsulation, was selected as a constant value and the quantity of lecithin was changed (table 1). Formulation number 2 (containing lecithin: drug: cholesterol ratio of 100: 10: 10) having about 90.05% encapsulation was selected as the best formulation ($P < 0.05$). Results of particle size determination showed 50% of MLV and SUV liposomes had diameter below 33.80 μm and 22.09 μm , respectively. According to the results of particle size, sonication is an important parameter in preparation of SUV liposomes. Also the results of microscopy indicated the SUV liposomes may be obtained by sonication of MLV liposome in a bath sonicator (fig 1a and 1b).

It has been previously reported that sonication time is an important parameter in the liposome preparation, and also increasing the time of sonication is necessary to decrease the polydispersity [13]. Also, loading of essential oil of *Eucalyptus camaldulensis* in liposomes, showed that homogenization is another factor that can significantly affected the liposomal particle size [14].

According to the results, release of drug from all of the formulations followed zero- order kinetics. The results showed that the release from SUV liposomal gel has the most regular and showed the least interaction between the drug and polymers (table 3, fig 2). The drug release from hydroalcoholic gel had the most irregularity and the highest interaction (table 3, fig 2). Also similar results obtained from MLV liposomal gel and the blank formulation indicated that a high percentage of the drug may be attached to external layer of liposomes (table 3, fig 2).




Fig 1: a) Image of MLV liposomes of Acitrecin and Methotrexate b) Image of SUV liposomes of Acitrecin and Methotrexate (200 X).

The stability of the vesicle dispersions was monitored for 3 months and the results indicated that there was no significant change in the appearance and color of the samples during the storage.

Acitrecin and Methotrexate is a glucocorticoid and is useful for treatment of inflammatory diseases. However, there is a tendency to reduce its use due to its complicated pharmacokinetic characteristics [11]. Many studies have been reported for improvement of tissue distribution and targeting of glucocorticoids drugs by using nano drug delivery system such as liposome and solid lipid nanoparticle (SLN). Ghanshyam *et al* in 2011 have been investigated the formulation and characterization SLNs dry powder inhaler containing Acitrecin and Methotrexate and the results showed SLNs are suitable carriers for incorporating the drug [11].

Table 3: The percentage release of different formulation gel of Acitrecin and Methotrexate.

Time (min)	Release of hydroalcoholic gel (%)	Release of MLV liposomal gel (%)	Release of SUV liposomal gel (%)	Release of blank MLV liposome gel plus the drug (%)
15	0.68±0.56	5.81±0.84	5.81± 4.78	6.66±0.97
30	4.75±2.14	8.99±0.27	9.85±4.68	7.83±1.03
45	6.91±2.56	12.43±0.58	12.66±4.45	9.59±1.24
60	9.49±1.83	14.99±0.77	19.16±3.30	14.59±2.62
90	13.22±1.63	21.12±1.09	27.58±3.47	19.20±2.30
120	17.49±1.59	27.51±1.03	37.33±3.09	25.53±2.57
150	22.75±1.56	33.69±1.66	44.00±4.54	31.77±3.39
180	25.84±1.79	39.69±1.66	53.70±6.52	38.59±3.63
210	31.20±2.91	45.26±1.90	61.39±8.47	45.37±3.68
240	37.38±2.71	51.90±1.70	69.49±11.20	53.71±4.90
270	42.40±2.20	58.59±2.07	76.72±11.29	60.86±5.84
300	48.15±1.49	69.91±2.38	83.90±11.17	67.87±5.82




Fig 2: The percentage release of Acitrecin and Methotrexate from different gel formulations

Topical application of liposome vesicles has many advantages over the conventional dosage forms. It has been suggested that the formulated liposomes may be applied to the skin as gel. It has been previously showed that the liposomal gel of Lidocaine HCL may perform therapeutically better effects than the conventional formulation [15]. Samani *et al* in 2009 showed that cyproterone acetate containing liposomes had $74 \pm 6.11\%$ loading efficiency and the liposomal formulation has better penetration potential than conventional cyproterone acetate formulation [7]. Agarwal *et al* in 2002 have evaluated the miconazol (MCZ) nitrate loaded topical liposome *in vitro*. The results have shown the entrapment of MCZ in liposome can facilitate localized delivery of the drug [10]. Also, formulation of celecoxib loaded liposomes and the study of their release behavior indicated that the potential advantages of celecoxib loaded liposome as carrier system [13].

CONCLUSION

According the results of the present study, incorporation of Acitrecin and Methotrexate in liposome in proper amounts and incorporation of liposomes in gel formulation has many advantages over the conventional dosage forms, including prolonged and controlled release topical dosage delivery.

REFERENCE

- [1] Chetanachan P, Akarachalanon P, Worawirunwong D, Dararutana P, Bangtrakulonoth A, Bunjop Kongmuang S. *Advanced Materials Research* 2008;55-57:709-711.
- [2] Maghaby G, Barry BW, Willians AC. *European J Pharmaceutical Sciences* 2008; 34: 203- 222.
- [3] Egbaria K, Weiner N. *Advanced drug delivery reviews* 1990; 5: 287-300.
- [4] Sharma VK, Mishra DN, Sharma AK, Srivastava B. *IJCPR* 2010; 1(2): 6-16.
- [5] Goyal P, Goyal K, Kumar SGV, Singh A, Katare OP, Mishra DN. *Acta Pharm* 2005; 55: 1– 25.
- [6] Shazly G, Nawroth T, Langguth P. *Dissolution Technologies* 2008; 7-10.

- [7] Mohammadi Samani S, Montaseri H, Jamshidnejad M. Iranian J of Pharmaceutical Sciences 2009; 5(4): 199-204.
- [8] Saibaba, S.V.; Kumar, M.S.; Ramu, B. Pharmaceutical impurities and their characterization. European J Pharm Med Res 2016, 3(5), 190-196.
- [9] Ortan A, Campeanu GH, Pirvu CD, Popescu L. Roumanian Biotechnological letters 2009; 14(3): 4411-4417.
- [10] Agarwal R, Katare OP. Pharmaceutical Technology 2002; 48-60.
- [11] Umaratiya Ghanshyan M, Patel Priyal R, Patel Javadan K. IJPRS 2011; 1(3): 662-673.
- [12] Dabbagh MA, Ford JL, Rubinstein MH, Hogan JE, Rajabi Siahboomi AR. Pharmaceutical Development and Technology 1999; 4(3): 136-142.
- [13] Moghimipour E, Handali S. Advanced Pharmaceutical Bulletin 2012; 2(1): 93-98.
- [14] Moghimipour E, Aghel N, Zarei Mahmoudabadi , Ramezani Z, Handali S. Jundishapur J Nat Pharm Prod 2012; 7(3): 117-122.
- [15] Glavas Dodov M, Simonoska M, Goracinova K. Bulletin of the Chemists and Technology of Macedonia 2005; 24(1): 59-65.

