

STANDARDISATION OF POLYHERBAL GEL FOR REJUVENATION OF HAIR GROWTH

Hanwathe Parameshwar*,A.V.Jithan,Andhe Balalji ,Kosna Randheer Reddy ,
Bonkuri Kumar ,P.Pravalika,Thatikanti Sushmitha.
Omega College Of Pharmacy,Edulabad,Ghatkesar, Hyderabad.

Abstract

The present study was designed to formulate and evaluate polyherbal gel containing extracts of Lagenaria siceraria, Trichosanthis cucumarina ,Tridax procumbens . Preliminary phytochemical screening were carried out for all the plants and its extracts to determine the presence of active principle in plants.Poly herbal gel was prepared with water soluble polymer Carbopol, propylene glycol400, povidone, triethanolamine to bring a good absorption capacity of the plant extracts on scalp. The standardization parameters of the gel are viscosity, pH, Homogeneity, Spreadability, content uniformity, skin irritation test all were carried out to bring a quality, purity and safety of the prepared gel formulation with the reference who applied gel without the extract. The growth of hair measured by trichoscope and the growth was completely observed after the 90 days Hence, from these studies it is concluded that the prepared poly herbal gel containing Lagenaria siceraria, Trichosanthis cucumarina, Tridax procumbens proved hair growth activity

Keywords: Carbopol 934, Hair growth initiation, Hair growth completion, Histopathology, Hair follicles

Introduction

Recently, the number of men and women who suffered from hair loss and/or hair thinning is increasing in worldwide. Hair loss is a dermatological disorder, and the surge for discovering natural products with hair growth promoting potential is continuous [1,2]. Hair loss or alopecia is a common patient complaint and a source of significant psychological and physical distress [3]. Many factors such as metabolism, hormones, heredity and side effects of antineoplastic and immunosuppressant drugs, have been negatively affecting the healthy growth of hair. According to one survey, androgenic alopecia on its own eventually affects approximately 50% of the world's adult population [4,5]. In androgenic alopecia it is assumed that the genetically predisposed hair follicles are the target for androgen - stimulated hair follicle miniaturization, leading to gradual replacement of large, pigmented hairs (terminal hairs) by barely visible, depigmented hairs (vellus hairs) in affected areas [4]. It is dihydrotestosterone mediated process, characterized by continuous miniaturization of androgen reactive hair follicles and accompanied by per follicular fibrosis of follicular units in histological examination [6-9].

Lagenaria siceraria is a small shrub under cucurbitaceae family. It is grow over the tropical and subtropical area and easily available. *Lagenaria siceraria* has various ethno botanical applications and medicinal claims. Its fruits have been use vegetables and the ethnomedicinal uses of leaves shows hair growth activity *Trichosanthes cucumerina* , is a small shrub under cucurbitaceae family. It is grow over the tropical and subtropical area and easily available. Its ethnomedicinal uses shows action on hair growth.

Tridax procumbens is a small weeds procumbent herb, Asteraceae family and easily available through the tropical and subtropical areas. Its leaves ethnomedical uses claim to have hair restoring properties. Among topical formulation, the gel formulation is more suitable for topical application and produce cooling effects.

PREPARATION OF EXTRACT

Extraction is the preliminary step involved in the phytochemical studies. Based on solvent's polarity metabolites are extracted and according to the solubility of the constituents in the solvent. The method of extraction is hot percolation method.

METHODOLOGY

LIST OF MATERIALS AND THEIR USES IN FORMULATION

The materials used in the study were as follows:

Table 1 List of materials and their uses in formulation

S.no	Name of the Materials	Use in formulation
1.	<i>Legenaria siceraria</i> (Leaves)	Active Ingredient
2.	<i>Trichosanthes cucumerina</i> (Leaves)	Active Ingredient
3.	<i>Tridax procumbens</i> (Leaves)	Active Ingredient
4.	Carbopol 934	Gelling agent
5.	Povidone	Suspending agent
6.	Polyethylene Glycol	Moisturizing agent
7.	Tuber rose oil	Flavoring agent
8.	Demineralized Water	Solvent

❖ HOT PERCOLATION METHOD

About 200g of coarsely powdered plant was extracted with solvents of increasing polarity like Hexane, Chloroform, Ethyl acetate and Ethanol at 60-70°C. Each extract was concentrated using rotary vacuum evaporator. The percentage yield, color and consistency of all the extracts were noted and were taken up for further

QUALITATIVE PHYTOCHEMICAL ANALYSIS

Qualitative analysis for various phytoconstituents in the dried powders and extracts all the raw materials were carried out using different reagents are mentioned below.

DEVELOPMENT AND STANDARDISATION OF POLY HERBAL FORMULATION

The prepared extracts were taken for the preparation of topical gel with water soluble gelling agent Carbopol 934, Propylene glycol 400, Povidone, Glycerin, Distilled water.

Method for Preparation of Gel Containing Extracts (1% w/w)

1 g of Carbopol 934 was dispersed in 50 ml of hot distilled water with continuous stirring. 5ml of distilled water was taken and required quantity of povidone was dissolved. Cool the solution, then to that added Propylene glycol 400.

Further required quantity of prepared leaves extracts were mixed to the above mixture and volume made up to 100 ml by adding remaining distilled water. Finally full mixed ingredients were mixed properly to the Carbopol 934 gel with continuous stirring and tri-ethanolamine was added drop wise to the formulation for adjustment of required skin pH (6.8-7) and to obtain the gel at required consistency.

Table .2 Formula for gel preparation

Ingredients	Quantity Specified
Carbopol-934	1 g
Propylene glycol-400	Quantity sufficient
Povidone	5%
Triethanolamine	1.2ml(quantity sufficient)
Distilled water	Make up to 100

Standardisation of Topical Gel Formulation

- **Physical Evaluation**

Physical parameter(50)s such as colour and appearance were checked.

- **Measurement of pH**

pH of the gel was measured by using pH meter.

- **Spreadability**

Spreadability of the prepared gel was determined by wooden block and glass slide apparatus.

Apparatus consists of a wooden block, which was provided by a pulley at one end. Spreadability was measured on the basis of slip and drag characteristics of gels. An excess of gel (about 2g) under study was placed on this ground slide. The gel was then placed between this slide and another glass slide having the dimension of fixed ground slide and provided with the hook. A 1 kg weighted was placed on the top of the two slides for 5 minutes to expel air and to provide a uniform film of the gel between the slides. Excess of the gel was scrapped off from the edges. The top plate was then subjected to pull of 80gms. With the help of string attached to the hook and the time (in seconds) required by the top slide to cover a distance of 7.5 cm be noted. A shorter interval indicates better Spreadability.

Spreadability was calculated using the following formula:

$$S = M \times L / T$$

Where, S = Spreadability,

M = Weight in the pan (tied to the upper slide), L =

Length moved by the glass slide and

T = Time (in sec.) taken to separate the slide completely each other.

Viscosity

Viscosity of gel was measured by using Brookfield viscometer with spindle.

Homogeneity

All developed gels were tested for homogeneity by visual inspection after the gels have been set in the container. They were tested for their appearance and presence of any aggregates.

Skin irritation test

Test for irritation was performed on human volunteers. The prepared gel was applied

on area of 2 square inch to the back of hand to volunteers who are selected for the evaluation

of hair growth gel. The volunteers were observed for lesions or irritation.

CLINICAL EVALUATION OF HAIR GROWTH ,

The prepared and standardized gel formulation was evaluated clinically. The clinical evaluation carried out in the Department of Dermatology, Government General Hospital, Hyderabad.

Study groups and duration of study

30 volunteers selected according to the inclusion criteria. They are to be divided into 2 groups control and test. The control group receives gel without extract medication. The test group receives the formulated herbal gel. Duration of the study is 90days.

Application of gel

The formulated gel is applied over the scalp during the course of study.

If there is any sign of irritation or discomfort to the patient, the gel is discontinued.

Evaluation parameters for hair growth (During the period of study)

- Measurement of hair texture, examined by simple method of touching the patient's hair,
- Hair density/cm.sq area :examined at a fixed site of scalp with Trichoscope
- Hair loss: counted after constant combing of patient's hair for one minute with the comb

RESULTS AND DISCUSSION

PREPARATION OF EXTRACTS

The shade dried all plants materials raw were extracted in soxhlet extractor with the universal solvent ethanol. All the extracts were concentrated using rotary vacuum evaporator. The percentage yield was calculated for every extract in terms of dried weight of plant materials. The colour and consistency of the concentrated extracts are given in table no.3.

QUALITATIVE ESTIMATION OF PHYTOCONSTITUENTS

The raw material powders and all the extracts were subjected to qualitative phytochemical analysis to identify the various phytoconstituents present in it, as per the standard procedures. The results are given in the Table 8.8

FLUORESCENCE ANALYSIS

Fluorescence analysis for the extracts and the powdered drug were carried out with various reagents to identify the presence of chromophores. The importance of fluorescence analysis is that UV light shows the fluorescent nature of the compound whereas fluorescence cannot be observed in day light. Hence it is performed according to the standard procedures. No fluorescence was observed for the powder as well as extracts indicating the absence of chromophore in the plant.

Table 3. Percentage yield of extracts

S.NO	PLANT NAME	SOLVENT	METHOD OF EXTRACTION	PHYSICAL NATURE	COLOUR	YIELD (%W/W)
1.	<i>Legenaria siceraria</i> (Leaves)	Ethanol (95%)	Continuous Hot percolation method using Soxhlet apparatus	Semisolid	Dark green	5.58
2.	<i>Trichosanthes cucumerina</i> (Leaves)			Semisolid	Dark green	4.76
3.	<i>Tridax procumbens</i> (Leaves)			Semisolid	Dark green	6.46

Table 4. Preliminary phytochemical analysis of powder and extracts of raw materials

Chemical constituents	<i>Lagenaria siceraria</i>		<i>Trichosanthes cucumerina</i>		<i>Tridax procumbens</i>	
	Powder	Extract	Powder	Extract	Powder	Extract
Steroids	+	+	+	+	-	-
Glycosides	-	-	-	-	+	+
Saponins	-	-	-	-	-	-

Flavonoids	+	+	+	+	+	+
Tannins	+	+	+	+	+	+
Proteins	+	+	+	+	-	-
Alkaloids	-	-	-	-	-	-
Carbohydrates	+	+	+	+	+	+
Terpenoids	-	-	-	-	-	-
Fats and oils	-	-	+	+	-	-

+ indicates presence , - indicates absence

QUANTITATIVE ESTIMATION

❖ TOTAL FLAVONOID CONTENT

The determination of total flavonoids from various extracts of *Lagenaria siceraria*, *Trichosanthes cucumerina*, *tridax procumbens* were performed with the Quercetin standard. The accuracy of test made by the serial dilution of Standard and the absorbance was measured spectrophotometrically at 435 nm. The obtained data were plotted as a Standard Calibration curve (Fig.1).

Standard Calibration Curve for Quercetin

To determine the accuracy of the flavonoid compound by plotted the standard absorbance obtained from spectrophotometrically. The calibration curve done by made serial dilution (20mcg, 40mcg, 60mcg, 80 mcg, 100 mcg) of quercetin Standard stock solution, the absorbance plotted against concentration

Table 5.: Spectrophotometric absorbance of Standard and Sample

S.No	Concentration of standard solution($\mu\text{g/ml}$)	Absorbance(435nm)
1.	20	0.135
2.	40	0.175
3.	60	0.213
4.	80	0.306
5.	100	0.347
6.	<i>Lagenaria siceraria</i>	0.178
7.	<i>Trichosanthis cucumarina</i>	0.165


Table 6 . Percentage yield of total flavonoid

S.NO	Sample	Concentration Obtained (mg/gm)	Percentage of Flavonoids Present
1.	<i>Lagenaria siceraria</i>	43.43	4.3
2.	<i>Trichosanthis cucumarina</i>	39.85	4
3.	<i>Tridax procumbens</i>	33.72	3.4

❖ TOTAL PHENOLIC CONTENT

Total phenolic content of the individual extract was determined and compared with that of standard.

Figure .1: Standard Calibration Curve for Gallic acid



(Table 7)

Table 7. Percentage of phenolic content present in each extract

S.NO	Sample	Concentration Obtained (mg/gm)	Percentage of phenolic content Present(%w/w)
1.	<i>Lagenaria siceraria</i>	204.33	20.4
2.	<i>Trichosanthis cucumarina</i>	168.27	16.8

Table 9. STANDARDIZATION OF PREPARED POLY HERBAL GEL

S. No.	Parameters	Results
1.	Physical appearance	Light green
2.	pH	7.1

3.	Spreadability	Good
4.	Viscosity	43560 cps
5.	Homogeneity	Excellent
6.	Skin irritation test	No irritation

CLINICAL EVALUATION OF HAIR GROWTH

The prepared and standardized ploy herbal gel formulation was tested on human volunteers. Before the clinical evaluation was carried out ethical clearance was approved by Institutional Ethical Committee. The hair length and diameter was measured by trichoscope and the results were tabulated. Every testing interval the patients scalp photos taken with the trichoscope (Fig 3, Fig. 4).The number of hair present in scalp was counter as per square area. The obtained mean value from hair count in 1 cm square area for test group was plotted on graph


Figure .3 Hair in Scalp



Figure .4 Hair in Scalp



Fig..5 Mean No. of Hair count



Mean No. of hair count in/cm. sq. area for Test group

Fig .6 Mean No. of hair count For Control Group




Table 10 . Meant value of hair count

No. of Days	Test Group	Control Group
0	12.07±2.6	11.4±2.3
15	12.07±2.8	11.3±2.03
30	15.5±2.6	12.2±2.3
60	19.5±2.8	13.2±2.4

90	25.9±5.0	13.3±2.4
----	----------	----------

Values are expressed as Mean ± SD, n=15

Table 11. Mean number of hair loss in Combing test

Parameter	Group	Initial 0 days	After 15 days	After 30 days	After 60 days	After 90 days
Mean hair loss (/combing)	Test	114.5± 6.4	80.4± 12.4	78.2± 11.4	69.3± 10.4	51.5± 15.4
	Control	113.2± 6.4	109.2± 6.4	103.4± 6.7	99.9± 7.9	95.2± 6.8
Perceptible Reduction in hairloss by Patients(%)	Test	-	34.1%	38.6%	45.2%	63.1%
	Control	-	4.0%	10.4%	13.7%	18.0%

The hair growth activity of the prepared gel formulation was tested on human volunteers and growth activity was compared with the control group. The obtained mean value (Table 10) from both test and standard group were compared Fig 7.

Fig..7Comparison of hair growth of Test with Control




Table 12 . Measurement of density

Group	Initial Mean density /cm.sq area (mm)	Mean density /cm.sq area After 15 days (mm)	Mean density /cm.sq area After 30 days (mm)	Mean density /cm.sq area After 60 days (mm)	Mean density /cm.sq area After 90days (mm)
Test	0.09±0.16	0.14±0.14	0.13±0.17	0.14±0.16	0.14±0.2

Control	0.08±0.17	0.07±0.16	0.08±0.18	0.11±0.10	0.11±0.14
----------------	-----------	-----------	-----------	-----------	-----------

CONCLUSION

Preliminary phytochemical screening were carried out for all the *Lagenaria siceraria*, *Trichosanthis cucumarina*, *Tridax procumbens* and their extracts to determine the presence of active principle in plants. Qualitative estimation of total flavonoid content and total Phenolic content were determined by spectrophotometrically all the extract showed significant amount of flavonoid and phenolic compounds. Poly herbal gel was prepared with water soluble polymer Carbopol, propylene glycol 400, povidone, triethanolamine to bring a good absorption capacity of the plant extracts on scalp.

The standardization parameters of the gel are viscosity, pH, Homogeneity, Spreadability, content uniformity, skin irritation test all were carried out to bring a quality, purity and safety of the prepared gel formulation. The prepared poly herbal formulation was taken for the determination of hair growth activity of the selected plants.

The clinical evaluation of prepared gel was carried on the human volunteers and compared with the reference who applied gel without the extract. The growth of hair measured by trichoscope and the growth was completely observed after the 90 days.

Hence, from these studies it is concluded that the prepared poly herbal gel containing *Lagenaria siceraria*, *Trichosanthis cucumarina*, *Tridax procumbens* proved hair growth activity.

REFERENCE

1. Victor R. Preedy. Handbook of hair in health and disease. In Wageningen Academic Publishers; 2012.
2. Ebling FJ. The biology of hair. *Dermatol Clin.* 1987;5(3):467–481.
3. Rhodes L, Harper J, Uno H, Gaito G, Audette-Arruda J, Kurata S, et al. The effects of finasteride (Proscar) on hair growth, hair cycle stage, and serum testosterone and dihydrotestosterone in adult male and female stump-tail macaques (*Macaca arctoides*). *J Clin Endocrinol Metab.* 1994 Oct;79(4):991–6.
4. D. Van Neste, J.M. Lachapelle, J.L. Antoine. Trends in Human Hair Growth and Alopecia Research. In: 1st ed. Netherlands: Kluwer Academic Publishers Group; 1989. p. 299–300.
5. Messenger AG, Slater DN, Bleehen SS. Alopecia areata: alterations in the hair growth cycle and correlation with the follicular pathology. *Br J Dermatol.* 1986;114(3):337–347.
6. Etefagh L, Nedorost S, Mirmirani P. Alopecia areata in a patient using infliximab: new insights into the role of tumor necrosis factor on human hair follicles. *Arch Dermatol.* 2004;140(8):1012–1012.
7. Ikeda T. A new classification of alopecia areata. *Dermatology.* 1965;131(6):421–445.
8. Jerry Shapiro, FRCPC. Hair Loss: Principles of Diagnosis and Management of Alopecia. In United Kingdom; 2002. p. 135–45.
9. Antonella Tosti, Bianca Maria Piraccini. DIAGNOSIS AND

- TREATMENT OF HAIR DISORDERS An Evidence Based Atlas. In United Kingdom: Taylor & Francis; p. 2006.
10. WILLIAM MONTAGNA, RICHARD A. ELLIS. In: The Biology of HAIR GROWTH. NEW YORK: ACADEMIC PRESS INC.; 1958. p. 173–85.
 11. Mecklenburg L, Tobin DJ, Müller-Röver S, Handjiski B, Wendt G, Peters EM, et al. Active hair growth (anagen) is associated with angiogenesis. *J Invest Dermatol.* 2000;114(5):909–916.
 12. Parasuraman S, Thing GS, Dhanaraj SA. Polyherbal formulation: Concept of ayurveda. *Pharmacogn Rev.* 2014;8(16):73–80.
 13. Petchi RR, Parasuraman S, Vijaya C. Antidiabetic and antihyperlipidemic effects of an ethanolic extract of the whole plant of *Tridax procumbens* (Linn.) in streptozotocin-induced diabetic rats. *J Basic Clin Pharm.* 2013 Sep;4(4):88–92.

