

A STUDY TO ASSESS THE ORIGIN, AND PATTERN OF IMPORTANT BLOOD VESSEL OF THE ABDOMINAL WALL.

Mr.Guntha Chinna Nagaraju, Research Scholar, Malwanchal University

Prof. Dr. Pawan Kumar ,Research Supervisor Malwanchal University

Introduction

Nature is a place where variations abound. However, even if many of the variants are minor, if they cause no disruption in the body's functions, they may be of great significance. There are developmental variations in the development of nerves, blood vessels, lymphatics, bones, joints, organs, and virtually all tissues in the human body. Anatomists and surgeons have long been fascinated by the interesting variances and anomalies that can be found in the origin and course of arterial branches. The inferior epigastric artery, also known as the deep inferior epigastric artery, is a blood vessel that supplies blood to the stomach.

As it is frequently referred to by surgeons, it is located bilaterally in the anterior abdomen wall of the thoracic cavity. It receives its supply from the external iliac artery, which is located in the extraperitoneal connective tissue immediately posterior to the inguinal ligament (inguinal ligament artery). It

In the following section, the tendon ascends in an oblique fashion along the medial edge of the deep inguinal ring.

Afterwards, it pierces the transversalis fascia and ascends in front of the arcuate scapula.

It continues upward between the rectus abdominis muscle and the posterior lamina of its sheath until it reaches the oblique abdominal muscles. It is divided into multiple branches, all of which are anastomosed with the branches of the superior epigastric artery (see figure).

The inferior epigastric artery has several branches, including the cremasteric, pubic, muscular, cutaneous, peritoneal, and anastomotic branches.

The public branch, in particular, is significant since it demonstrates a huge amount of dedication.

It is possible to have an obturator artery abnormality. The pubic branch of the obturator artery forms an anastomosis with this pubic branch. The foundation of this relationship has been laid.

Corona Mortis is the medical word for this condition (crown of death). It is possible that the crown of death will be lacerated if blind dissection is performed along the iliopectineal line throughout the procedure.

Orthopaedic surgeons, urologists, and gynaecologists are all concerned about the presence of aberrant vessels in the retropubic region, and they are not alone in their concern.

Furthermore, general surgeons are involved in this field of work through performing procedures.

Furthermore, the origin, branching pattern, and anastomosis pattern of the inferior epigastric artery are vulnerable to change.

In only 15 years, laparoscopic surgery has gone from being a rare and exceptional experience to being performed on an almost daily basis.

While the inferior epigastric artery and its branches are being inserted into the anterior abdominal wall during laparoscopic surgery, the inferior epigastric artery and its branches are at danger of injury.

As a result, it is necessary to investigate the anatomy of the inferior epigastric artery.

A lot of people are becoming interested in it.

Since the commencement of plastic surgery, the face has experienced significant transformations.

It has never failed to provide us with a reliable source of abundant, well-perfused tissue on a constant basis. The lower abdomen has been used in long-distance transfers after being used in pedicled and tubed flaps for a long period of time initially.

It was discovered that skin and fat were the most suitable materials for re-creating a woman's breasts.

The rectus abdominis musculocutaneous flap was the first of its kind to be utilised for the purpose for which it was intended.

Drever and Hartrampf demonstrated in 1977 and again a few years later that a skin island harvesting technique can be used to harvest the skin island transversely across the lower abdomen. It is oriented in a transverse direction.

By employing the rectus abdominis muscle (TRAM) flap, it was possible to reconstruct the breast using autogenous tissue.

It exhibited increased vascularity when compared to its pedicled cousin, owing to the robust blood supply given by the deep inferior epigastric system.

TRAM flaps can also be used to cover substantial flaws in the groyne, which is another application for the flaps.

As a result of the need for substantial groyne wounds as a result of trauma or cancer ablation, laparoscopic surgery has become a well-established diagnostic and therapeutic tool in the medical community. Large trocars must be placed through the anterior abdominal wall in order to perform many of these novel procedures. Because of the substantial vascular supply in the anterior abdominal region, these procedures increase the risk of injury to the vessels of the abdominal wall (the inferior and superior epigastric arteries) during surgery (Fig. 3). If the inferior epigastric artery is damaged, it can cause severe hypotension and even death if left untreated. If the anterior abdominal wall and its vascular architecture are well known, damage to the inferior epigastric artery during laparoscopic surgery is less likely to take place.

During femoral hernia treatment surgeries, the pubic branch of the inferior epigastric artery is in danger of being injured by the surgeon.

"Corona mortis" has been used to describe the connections between the pubic rami of the inferior epigastric artery (IEA) and the posterior obturator artery (OBA) for centuries (crown of death). Many procedures in the retropubic region necessitate the use of Corona Mortis treatment because it has the potential to produce serious bleeding.

It has been reported that the use of abdominal wall retention sutures can result in a pseudoneurysm of the inferior epigastric artery in some patients.

Historically, the inferior epigastric artery has been recommended for coronary bypass grafting because of its low thrombogenicity and low blood clotting time (CABG).

Ischemic necrosis is quite rare, and blood flow is good throughout the body.

compliance.

The blood supply to the breasts is provided by this inner artery, which branches out from the subclavian artery.

The superior epigastric artery, which connects the stomach to the inferior epigastric artery, is a blood vessel that runs through the stomach.

In the event of aorto-iliac occlusive disease, this acts as a critical collateral network for the patient's survival.

Erectile dysfunction is treated with the use of microvascular anastomosis.

This method makes use of the deep dorsal vein of the penis as well as the inferior epigastric artery to accomplish its goal.

The inferior epigastric artery serves as an interpositional conduit for grafting during the implantation process.

Restoration of arterial input to the transplanted liver is required when an acute hepatic artery blockage occurs during a liver transplantation procedure.

The inferior epigastric artery (IEA) is a good alternative for a graft because of its flexibility as a graft and rising recognition in the retropubic region.

The groin area, the anterior abdominal wall, and the inferior limb have all provided me with motivation to continue my research on the inferior limb and lower extremity.

The process that results in the formation of the inferior epigastric arterial trunk

a) The external iliac artery or any other circulatory system connection

a) The inguinal artery's position in relation to the origin of the external iliac artery In the rectus muscle, this is represented by the upper, middle, or lower third of the rectus muscle.

Before it enters the inferior epigastric artery, this artery can have either a single or a double stem.

A substance found in the rectus muscles

There are several branches of the inferior epigastric artery.

This condition is characterised by an irregularity in the route of the obturator artery and its connection to the anatomical ring of arteries that surrounds the lower leg.

In the inferior and superior epigastric arteries, there is an anastomosis.

The upper epigastric artery and the seventh epigastric artery are anastomosed at or above the umbilicus, depending on the location of the anastomosis.

The eighth factor is the distance between the inferior epigastric artery and key abdominal and pelvic landmarks.

The distance between the inferior epigastric blood vessel's midline and the inferior epigastric blood vessel's midline

"It's called the umbilicus."

Measure the distance between the inferior epigastric artery and the midline at the level of the inferior epigastric artery.

A symphysis is a connection between the umbilicus and the pubic symphysis.

When the inferior epigastric artery is measured at the level of the stomach, it is said to be

The symphysis pubis is a joint in the lower abdomen.

1) The distance between the tendon's origin and the point at which it makes contact with the The inferior epigastric artery connects the lateral rectus border to the inferior epigastric artery.

The inferior epigastric vein's anatomical length is measured in millimetres.

It passes through the rectus sheath from the time of its conception until the time of its entrance.

b) From the inferior epigastric artery's origin to the point of entry into the rectus muscle

The diameter of the inferior epigastric artery is

There are two ways to approach this situation.

b) At the rectus muscle's entry point into the inferior epigastric artery.

substance.

In this study, we've made a modest attempt to shed some light on the major artery of the abdomen wall, but we've come up short.

Methodology

The investigation was conducted in the form of a descriptive observational study. A total of 50 corpse samples were chosen for study (25 female and 35 male samples), with a fifth foetus cadaver being utilised as an additional reference. Medical colleges in the state of Maharashtra were chosen as the location.

We investigated the inferior epigastric arteries in fifty specimens, which included 50 adult human cadavers (25 males and 25 females) that had been kept in formalin for several months.

50 specimens were subjected to the conventional dissection method, predissectional dye injection, radiological study, and histological study procedures, all of which were performed. Predissectional redoxide with bull's fat mixture injection was performed on 50 specimens in order to analyse the branches of the inferior epigastric artery before dissection. A total of 5 foetuses (2 males and 3 females) were dissected and investigated as well.

Results

Several investigations into the inferior epigastric artery were carried out in order to determine the method of origin, the course of the artery, the branches, and the anastomoses that formed along the artery. At several points along the line from the midline to the inferior epigastric artery, measurements were collected. This should serve as a guideline for surgeons in order to perform a safe surgery while keeping the inferior epigastric artery free of obstruction. The inferior epigastric artery has been measured at two different levels, as has its diameter at two different levels. It would be beneficial to plastic surgeons who use the inferior epigastric artery as the pedicle to have a detailed study done on the inferior epigastric artery under the 10 parameters as described above.

According to the findings of the aforesaid authors, the inferior epigastric artery was shown to emanate from the external iliac artery in all 50 adult and 5 foetal specimens studied in the current investigation.

According to A.M. Buchanan (1906), the inferior epigastric artery originates from the medial side of the external iliac artery.

According to Donald Serafin (1996), the deep inferior epigastric artery develops from the medial aspect of the inferior epigastric artery. This is supported by other researchers.

Every one of the 50 adult specimens and five foetal specimens used in the current investigation showed that the inferior epigastric artery developed as a distinct trunk distinct from the external iliac artery. The present analysis did not uncover any shared trunks of origin with the obturator artery, medial circumflex femoral artery, or deep femoral artery, as previously reported.

Dschau (1936) described a case in which the inferior epigastric artery and the obturator artery were found to emerge as distinct arteries from the medial circumflex femoral artery.

It was discovered by Giovanni Teodori et al. (1984) that the inferior epigastric artery might come from the femoral or the deep femoral, all of which are below the inguinal ligament. The epigastric artery may develop from the obturator artery, which is supplied by the internal iliac, or it may be produced by two branches, one from the external iliac and the other from the internal iliac, or it may be formed by two branches from the external iliac and one from the internal iliac.

The inferior epigastric artery did not originate from the femoral, deep femoral, or medial circumflex femoral arteries or the obturator artery in any of the adult or foetal specimens used in the current study, nor did it originate from the obturator artery.

As previously demonstrated, none of the 50 adult specimens or five foetal specimens had the inferior epigastric artery created by two branches, one from the external iliac and the other from the internal iliac.

When 25 adult specimens were examined, the inferior epigastric artery was found to originate at the level of the inguinal ligament, as previously described by the authors of this study. The inferior epigastric artery arose at the level of the inguinal ligament in five of the five foetal specimens studied.

Following Morris (1893) and John E. Skandalakis (2004), it has been determined that the inferior epigastric artery originates immediately above the inguinal ligament.

According to the current study, the inferior epigastric artery arises above the inguinal ligament in 53% of adult cases, which is consistent with previous findings. The distance between the inferior epigastric artery and the inguinal ligament ranged between 0.5-2 cm, with an average of 1.1 cm, which is consistent with the findings of A.M. Buchanan, Sir John Bruce et al, Donald Serafin, and Michael S. Baggish and Mickey M. Karram, as well as other researchers.

In three foetal specimens, the inferior epigastric artery arose 0.4 cm above the inguinal ligament, indicating that it was a lateral branch. There is no mention of a foetal study in any of the authors' papers.

In the present investigation, the inferior epigastric artery was found to enter the rectus muscle in the lower third of 12 cases and in the upper third of 5 cases, which is more consistent with the findings of Milloy et al. in the previous study.

The inferior epigastric artery was found to enter the middle portion of the rectus abdominis in all of the foetal specimens studied. The authors of the above-mentioned paper make no mention of a study on foetuses.

Using 50 adult cadaveric halves, the present investigation discovered that the inferior epigastric artery had a single main stem in 83 percent of the specimens and a double main stem in 19 percent of the specimens, which is completely consistent with the author's findings.

All five of the foetal specimens studied had the inferior epigastric artery with a single stem before entering the rectus abdominis in all five of the foetal specimens.

In the present investigation of 50 adult specimens, the cremasteric, pubic, muscular, cutaneous, peritoneal, and anastomotic branches of the inferior epigastric artery were found in the present investigation, which corresponded to Buchanan's classification of inferior epigastric artery branches.

The cremasteric branch, pubic branch, and muscular branches were the only ones that could be found in the foetal specimens.

An investigation of 50 adult cadaveric halves revealed that in eight specimens, the pubic branch of the inferior epigastric arterial was enlarging, and this artery was replaced by and flowed through the obturator canal as the anomalous obturator arterial in the present study.

The pubic branch of the inferior epigastric artery was found to be enlarged in the current investigation, and it took the place of the obturator artery as the aberrant obturator artery. Suppa, Pungapong, and SathonThum findings Umnausuk's findings that the aberrant obturator artery occurs at a rate of 18 percent in the present study are the closest to the findings of J.L. Braithwaite and Suppa ut Pungapong and SathonThum-Umnausuk.

According to Henry Gray (1858), the aberrant obturator artery is located near the medial border of the femoral ring and is a source of pain.

In the present investigation of 50 adult cadaveric halves, the aberrant obturator artery was found to be associated to the medial border of the femoral ring in 18.7 percent of the instances, which is the percentage that is closest to the values of A Lee McGregor.

Last (1954) hypothesised that the aberrant obturator artery could be located on the lateral border of the femoral neck. Is this correct?

In the current study, the aberrant obturator artery descended on the lateral boundary of the femoral ring in 82.1 percent of patients, which is consistent with the findings of A. Lee McGregor's research (1975).

When Thomas Dwght and colleagues (1907) investigated aberrant obturator artery degeneration, they discovered it in 27.5 percent of cases.

The aberrant obturator artery lying across the femoral ring was not observed in any of the 50 adult specimens examined in this investigation.

Anastomoses were detected in 14 adult specimens (26 percent) in the current investigation, while no anastomoses were found in 77 percent of the specimens.

Among the cases studied by Milloy et al. (1960), one anastomosis was found between the superior epigastric artery and the inferior epigastric artery in 18.3 percent of the cases.

Multiple anastomoses were found in 8% of the specimens in the current investigation of 50 adult specimens, which is similar to the finding of Milloy et al. in their analysis of the same population.

In any of the five foetal specimens studied, there were no anastomoses found between the inferior epigastric artery and the superior epigastric artery. The authors of the above-mentioned paper make no mention of foetal research.

Anastomoses at or near the level of the umbilicus were not discovered in any of the adult specimens examined in this investigation.

Anastomoses between the superior and inferior epigastric arteries were absent from any of the five foetal tissues examined. One or more of the authors have made no mention of the foetal study.

The distance between the inferior epigastric artery and the midline at the level midway between the umbilicus and the pubic symphysis in the present investigation of 50 adult cadaveric halves was 3.7 + 0.3 cm on the right and 3.6 + 0.2 cm on the left, which does not agree with the author's findings.

Approximately 6 – 7 cm separates the midline (which is located above the upper edge of the symphysis pubis) from the inferior epigastric arteries, according to Mickey M. Karram (2001).

It was discovered in this investigation that the average diameter of the inferior epigastric artery in 50 adult cadaveric halves was 2.3mm, which is comparable to the value reported by Kim D I et al (2003).

Histology

In this investigation, the intima of the inferior epigastric artery was found to be thinner than the intima of the internal thoracic artery, which was consistent with the findings of A. Wahba and colleagues (1994). In this investigation, the media and intima thicknesses were found to be lower in the inferior epigastric artery than in the internal thoracic artery in this investigation, which matched the author's findings. In the current study, the internal thoracic artery was shown to have more elastic fibres in the media, while the inferior epigastric artery had more smooth muscle fibres, similar to the findings of A. Wahba et al.

Conclusion

The inferior epigastric artery, the most significant and largest blood vessel of the anterior abdominal wall, was investigated in great detail using conventional dissection, predissectional dye injection, angiographic, and histological studies. The study's conclusions were backed up by earlier research. Bringing together a comprehensive study of the inferior epigastric artery's origin and course, branches and anastomosis, length, diameter, and distance from important abdominal wall landmarks under one roof will be of great benefit to plastic surgeons who consider lower abdomen skin and fat to be an ideal material for breast reconstruction, and cardiothoracic surgeons who are evaluating the inferior epigastric artery as a potential alternative conduit for coronary artery bypass surgery..

Reference

1. Masia J, Larrañaga J, Clavero JA et al The value of the multidetector row computed tomography for the preoperative planning of deep inferior epigastric artery perforator flap: our experience in 162 cases. *Ann Plast Surg* 2008; 60: 29–36 [PubMed] [Google Scholar]
2. Smit JM, Dimopoulou A, Liss A et al Preoperative CT angiography reduces surgery time in perforator flap reconstruction. *J Plast Reconstr Aesthet Surg* 2009; 62: 1,112–1,117 [PubMed] [Google Scholar]
3. Ghattaura A, Henton J, Jallali N et al One hundred cases of abdominal-based free flaps in breast reconstruction. The impact of preoperative computed tomographic angiography. *J Plast Reconstr Aesthet Surg* 2010; 63: 1,597–1,601 [PubMed] [Google Scholar]

4. Rozen WM, Phillips TJ, Ashton MW et al Preoperative imaging for DIEA perforator flaps: a comparative study of computed tomographic angiography and Doppler ultrasound. *Plast Reconstr Surg* 2008; 121: 9–16 [PubMed] [Google Scholar]
5. Moon HK, Taylor GI. The vascular anatomy of rectus abdominis musculocutaneous flaps based on the deep superior epigastric system. *Plast Reconstr Surg* 1988; 82: 815–832 [PubMed] [Google Scholar]
6. Rozen WM, Palmer KP, Suami H et al The DIEA branching pattern and its relationship to perforators: the importance of preoperative computed tomographic angiography for DIEA perforator flaps. *Plast Reconstr Surg* 2008; 121: 367–373 [PubMed] [Google Scholar]
7. Nahabedian MY, Momen B, Galdino G, Manson PN. Breast reconstruction with the free TRAM or DIEP flap: patient selection, choice of flap, and outcome. *Plast Reconstr Surg* 2002; 110: 466–475 [PubMed] [Google Scholar]
8. See MS, Pacifico MD, Harley O et al Incidence of ‘incidentalomas’ in over 100 consecutive CT angiograms for preoperative DIEP flap planning. *J Plast Reconstr Aesthet Surg* 2010; 63: 106–110 [PubMed] [Google Scholar]
9. Rozen WM, Ashton MW, Grinsell D et al Establishing the case for CT angiography in the preoperative imaging of abdominal wall perforators. *Microsurgery* 2008; 28: 306–313 [PubMed] [Google Scholar]
10. Rozen WM, Ashton MW, Grinsell D. The branching pattern of the deep inferior epigastric artery revisited in-vivo: a new classification based on CT angiography. *Clin Anat* 2010; 23: 87–92 [PubMed] [Google Scholar]
11. Pacifico MD, See MS, Cavale N et al Preoperative planning for DIEP breast reconstruction: early experience of the use of computerised tomography angiography with VoNavix 3D software for perforator navigation. *J Plast Reconstr Aesthet Surg* 2009; 62: 1,464–1,469 [PubMed] [Google Scholar]009.

